

Child Behavior Checklist
Grade 1 / Year 2
Fast Track Project Technical Report
Anne-Marie Iselin
05/06/2011

Table of Contents

- I. Scale Description
- II. Report Sample
- III. Scaling
- IV. Differences Between Groups
- V. Recommendations for Use

Citation

Instrument

Achenbach, T.M. (1991). Manual for the Child Behavior Checklist / 4 – 18 and 1991 Profile. Burlington, VT: University of Vermont Department of Psychiatry.

Report

Iselin, A. R. (2011). Child Behavior Checklist (Year 2) (Technical Report) [On-line]. Available: <http://www.fasttrackproject.org/>

I. Scale Description

The Child Behavior Checklist (CBCL) is composed of 113 items designed to record, in a standardized format, behavioral problems and competencies of children aged 4 through 18, as reported by their parents or other primary caregivers. The CBCL allows for the calculation of raw scores and t-scores, normed separately for girls and boys, in 8 different behavioral domains: Withdrawn, Somatic Complaints, Anxious/Depressed, Social Problems, Thought Problems, Attention Problems, Delinquent Behavior and Aggressive Behavior. The domain of Sex Problems (items 5, 59, 60, 73, 96, and 110) is *not* scored for this age group of children, although the items responses are available.

II. Report Sample

This technical report is based upon Year 2, Cohort 1 data, including both High-Risk Control and Low-Risk Normative samples. With missing cases excluded, the total sample size was 439 with 366 normative participants and 151 control participants. Seventy-eight children were part of both the normative group and the high risk control group. Our analyses however, examine non-overlapping samples such that the Normative sample is comprised of only Low-Risk respondents and the Control sample is comprised of only high risk respondents. Thus, the total sample size for our analyses is 439, with 285 Low-Risk Normative and 146 High-Risk Control subjects. More of the total sample was gathered from the Durham (28.5%) and Pennsylvania (27.6%) sites than from the Nashville (22.6%), and Seattle (21.4%) sites.

III. Scaling

A parent or primary caregiver rates each item on a 3-step response scale: (0) - not true, (1) somewhat or sometimes true and (2) - very true or often true. The item responses are scored using a program which accompanies the instrument, providing both raw total scores and t-scores, normed by gender, for each domain. The items comprising each behavioral domain are:

Withdrawn: 42, 65, 69, 75, 80, 88, 102, 103, and 111.

Somatic Complaints: 51, 54, 56a, 56b, 56c, 56d, 56e, 56f, 56g, and, if applicable, 56h.
 Anxious / Depressed: 12, 14, 31, 32, 33, 34, 35, 45, 50, 52, 71, 89, 103, and 112.
 Social Problems: 1, 11, 25, 38, 48, 55, 62, and 64.
 Thought Problems: 9, 40, 66, 70, 80, 84, and 85.
 Attention Problems: 1, 8, 10, 13, 17, 41, 45, 46, 61, 62, and 80.
 Delinquent Behavior: 26, 39, 43, 63, 67, 72, 81, 82, 90, 96, 101, 105, and 106.
 Aggressive Behavior: 3, 7, 16, 19, 20, 21, 22, 23, 27, 37, 57, 68, 74, 86, 87, 93, 94, 95, 97, and 104.

Additionally, the CBCL had three broad-band scores: Internalizing, Externalizing and Total Problems. The Internalizing scale is comprised of items from the Withdrawn, Somatic Complaints and Anxious / Depressed scales. The Externalizing scale is comprised of items from the Delinquent Behavior and Aggressive Behavior domains. The Total Problems scale is comprised of all items, except for items 2 and 4 (allergies and asthma).

IV. Differences Between Groups

A series of independent t-tests were conducted to assess differences between the Normative and High-Risk Control samples. For these analyses, the Normative sample consisted of only Low-Risk respondents. Results indicated statistically significant higher mean scores for the High-Risk Control sample across all domains and composite scores (i.e., broadband scores):

Low-Risk Normatives (n=285) vs High-Risk Controls (n=146)

Scales	t-value	df	p-value	Means (SD)	
				Low-Risk Normative	High-Risk Control
Withdrawn	3.47	255	<0.001	54.42(5.92)	56.75(6.93)
Somatic Complaints	2.27	270	0.02	56.15(6.68)	57.79(7.31)
Anxious / Depressed	3.60	255	<0.001	54.81(6.58)	57.51(7.72)
Social Problems	5.50	255	<0.001	55.32(7.22)	59.83(8.45)
Thought Problems	4.89	239	<0.001	54.12(6.07)	57.71(7.73)
Attention Problems	7.25	258	<0.001	55.14(6.73)	60.62(7.78)
Composites					
Delinquent Behavior	7.62	230	<0.001	55.72(6.57)	62.01(8.80)
Aggressive Behavior	7.43	239	<0.001	54.86(6.86)	61.03(8.74)
Internalizing	4.37	296	<0.001	52.01(10.03)	56.56(10.18)
Externalizing	9.52	310	<0.001	51.66(10.01)	60.99(9.40)
Total Problems	8.62	327	<0.001	52.25(10.58)	60.80(9.30)

* T-test with correction for degrees of freedom (df) given statistical inequality of variances.

V. Recommendations for Use

The CBCL provides a standardized, normative framework for evaluating a parent's or primary caregiver's observations of a child across 8 domains and 3 broadband scores, covering several DSM diagnoses. High-Risk control participants scored significantly higher than Low-Risk Control participants across all domains and composite scores.